

Positive Algebra

a collection of productive exercises

Sum and product(I)

$$\begin{array}{ccccccc} ? & + & ? & = & 12 \\ ? & \times & ? & = & ? \end{array}$$

Two natural numbers, you don't know which ones.

However, not everything is unknown!

This is what you know: if you **add** both numbers,
the result will be **12**.

Now both numbers are **multiplied**

◆ What can be the result ? Show all results you found.

Sum and product (II)

$$6 + 14 = 20$$

sum
of 6 and 14

$$6 \times 14 = 84$$

product
of 6 and 14

From two natural numbers you only know the **sum**, this is **20**.
If the numbers were 6 and 14 their **product** should be 84.
But there are other pairs of natural numbers which sum is equal to **20**.
Their product will differ from 84.

- ◆ Below you see a chart with the numbers 1, 2, ..., 100.
Colour all cells that correspond with a possible product.

1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20
21	22	23	24	25	26	27	28	29	30
31	32	33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48	49	50
51	52	53	54	55	56	57	58	59	60
61	62	63	64	65	66	67	68	69	70
71	72	73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88	89	90
91	92	93	94	95	96	97	98	99	100

- ◆ Write the results in a sequence from large to small.
Do you discover any pattern in this sequence? Which one?

Sum and product (III)

Two natural numbers have a **product** equal to **24**.

◆ What can be their **sum**?

Three natural numbers have a **sum** equal to **10**.

◆ What can be their **product**?

Sum and product (IV)

A and **B** represent natural numbers

$$A + B = 12$$



$$A \times B = \dots \text{ or } \dots \text{ or } \dots \text{ or } \dots \text{ or } \dots \text{ or } \dots \text{ or } \dots$$

$$A \times B = 12$$




$$A + B = \dots \text{ or } \dots \text{ or } \dots$$

$$A \times B \times C = 12$$




$$A + B + C = \dots \text{ or } \dots \text{ or } \dots \text{ or } \dots$$

Perimeter and area (I)



$$\begin{aligned} \text{perimeter} &= L + W + L + W \\ &= 2 \times (L + W) \\ &= 2 \times L + 2 \times W \end{aligned}$$



$$\text{area} = L \times W$$

The length and width of a rectangle are a whole number of centimeters, but you don't know how many. The **perimeter** is equal to 18 centimeters.

- ◆ Draw all possible rectangles.

How many square centimeters can be the **area**?

The length and width of a rectangle are a whole number of centimeters.

You have to know that the **area** is equal to 18 cm.


- ◆ Draw all possible rectangles.

How many centimeters can be the **perimeter**?


A rectangle has **perimeter** of 22 cm and **area** 28 cm².

- ◆ How long and how wide is the rectangle?

Perimeter and area (II)



$$\text{perimeter} = 4 \times A + 4 \times B$$



$$\text{area} = 3 \times A \times B$$

A and B represent whole numbers.



Suppose the **perimeter** of the shape above is 52 cm.

◆ What can be the **area**?

Suppose the **area** of the shape above is 42 cm².



◆ What can be the **perimeter**?

Perimeter and area (III)







- ◆ Fill in expressions for **perimeter** and **area**.
- ◆ Design a problem about perimeter and area of such a cross figure.


Find the three numbers




◆ Fill in the missing numbers:





How old?



How heavy?



How expensive?




Dot patterns (I)

Groups of birds sometimes fly in a **V-pattern**.




V-pattern with dots:



The figure shows the first four V-patterns. Each pattern has a number in the sequence. Below you see a V-pattern with 17 dots.


◆ Which number in the sequence has this V-pattern?




- ◆ How many dots has the V-pattern with number 85 in the sequence?
- ◆ Does exist a V-pattern with 35778 dots? Why or why not?
- ◆ Give a rule to find the number of dots of a V-pattern, knowing the number in the sequence.
- ◆ Represent this rule by a direct formula; use the letters n and V (n = number in the sequence, V = number of dots)

Dot patterns (II)

During a show a squadron of airplanes flew in a W-formation.



Look at the begin of a sequence of W-patterns:



◆ Fill in the table:

<i>pattern number</i>	1	2	3	4	5	6	
<i>number of dots</i>	5	



- ◆ How many dots are in the W-pattern number 25?
- ◆ Find a direct formula to describe the number of dots in any W-pattern.
(n = number in the sequence, W = number of dots)
- ◆ What is the relationship between a W-number and a V-number corresponding with the same number n ?
Is this $W = 2 \times V$ or not? Explain your answer
- ◆ Choose another letter-pattern by yourself and find a corresponding formula.

Dice patterns



Of course you know the six dot patterns on a dice:





- ◆ Which pattern, **a** or **b**, has the biggest number of dots?
You can answer this question without counting the dots!





- ◆ Which pattern, **c** or **d**, has the biggest number of dots?:



Operating with number strips (I)



Operating with number strips (II)



Operating with number strips (III)

$$5 \times \begin{array}{|c|} \hline 1 \\ \hline 3 \\ \hline 5 \\ \hline 7 \\ \hline 9 \\ \hline 11 \\ \hline 13 \\ \hline \end{array} = \begin{array}{|c|} \hline \\ \hline \\ \hline \\ \hline \\ \hline \\ \hline \\ \hline \\ \hline \end{array}$$

$$5 \times \boxed{} = \boxed{}$$

$$\frac{1}{2} \times \begin{array}{|c|} \hline 10 \\ \hline 16 \\ \hline 22 \\ \hline 28 \\ \hline 34 \\ \hline 40 \\ \hline 46 \\ \hline \end{array} = \begin{array}{|c|} \hline \\ \hline \\ \hline \\ \hline \\ \hline \\ \hline \\ \hline \end{array}$$

$$\frac{1}{2} \times \boxed{} = \boxed{}$$

Operating with number strips (IV)


$$\begin{array}{c}
 \text{3} \times \\
 \begin{array}{|c|}
 \hline
 2 \\
 \hline
 3 \\
 \hline
 4 \\
 \hline
 5 \\
 \hline
 6 \\
 \hline
 7 \\
 \hline
 8 \\
 \hline
 \end{array}
 \end{array}
 +
 \begin{array}{c}
 \text{2} \times \\
 \begin{array}{|c|}
 \hline
 1 \\
 \hline
 3 \\
 \hline
 5 \\
 \hline
 7 \\
 \hline
 9 \\
 \hline
 11 \\
 \hline
 13 \\
 \hline
 \end{array}
 \end{array}
 =
 \begin{array}{|c|}
 \hline
 \\
 \hline
 \\
 \hline
 \\
 \hline
 \\
 \hline
 \\
 \hline
 \\
 \hline
 \\
 \hline
 \\
 \hline
 \end{array}$$

$$\text{3} \times \boxed{} + \text{2} \times \boxed{} = \boxed{}$$

$$\begin{array}{c}
 \text{2} \times \\
 \begin{array}{|c|}
 \hline
 3 \\
 \hline
 8 \\
 \hline
 13 \\
 \hline
 18 \\
 \hline
 23 \\
 \hline
 28 \\
 \hline
 33 \\
 \hline
 \end{array}
 \end{array}
 -
 \begin{array}{c}
 \text{5} \times \\
 \begin{array}{|c|}
 \hline
 1 \\
 \hline
 3 \\
 \hline
 5 \\
 \hline
 7 \\
 \hline
 9 \\
 \hline
 11 \\
 \hline
 13 \\
 \hline
 \end{array}
 \end{array}
 =
 \begin{array}{|c|}
 \hline
 \\
 \hline
 \\
 \hline
 \\
 \hline
 \\
 \hline
 \\
 \hline
 \\
 \hline
 \\
 \hline
 \end{array}$$

$$\text{2} \times \boxed{} - \text{5} \times \boxed{} = \boxed{}$$

Strips and charts (I)




- ◆ Find the number that corresponds with $m = 3$ and $n = 2$
- ◆ Also for $m = 3$ and $n = 5$

- ◆ A horizontal row in the chart corresponds to $m = 4$. Which one?
- ◆ Arow in the chart corresponds to $n = 5$. Which one?


- ◆ Which numbers from the chart correspond to $m = n$?
- ◆ Make a number strip for these numbers with the corresponding expression.

Strips and charts (II)





- ◆ Fill in the chart. Which expression fits with the chart?
- ◆ Which strip fits with $m = 3$? What is the corresponding expression?
- ◆ Same questions for $n = 0$.
- ◆ Also for $m = n$.
- ◆ Also for $m = n + 1$.

Strips and charts (III)





- ◆ Which expression fits with the chart?
- ◆ From which strips can the chart be made?

Operating with expressions (I)







Operating with expressions (II)



Vertices, edges, faces (I)

The beginning of a sequence of **pyramids**.

- Explain the numbers below the pyramids.



For any pyramid we call:

V = the number of vertices,

E = the number of edges ,

F = the number of faces.

Example: for a quadrilateral pyramid you have $V = 5$, $E = 8$, $F = 5$





- Fill in the missing numbers:

n	V	E	F
3			
4	5	8	5
5			
6			
7			

- Find direct formulas for V , E and F related to n .
- The four pyramids above satisfy the relation: $V + F = E + 2$. Check this.
- Explain that $V + F = E + 2$ is true for every pyramid, using the direct formulas for V , E and F .

Vertices, edges, faces (II)

Here is a sequence of **prisms**.



Consider the numbers of vertices, edges and faces.

- Fill in the missing numbers:

n	H	R	V
3			
4	8	12	6
5			
6			
7			


- Now find direct formulas for V , E and F related to n :

$$H = \dots\dots\dots \quad R = \dots\dots\dots \quad V = \dots\dots\dots$$

- Does the equality $V + F = E + 2$ valid for each prism?
Use the direct formulae for V , E and F to investigate this.

Vertices, edges, faces (III)

Here is a sequence of **towers**:



Consider again the numbers of vertices, edges and faces .

- Fill in:


n	H	R	V
3			
4	9	16	9
5			
6			
7			

- Find direct formulas for V , E and F of a n -sides tower.

$$V = \dots\dots\dots E = \dots\dots\dots F = \dots\dots\dots$$






- For each tower valids: $V + F = E + 2$.
How can you explain this?

Formulas for perimeters (I)




$$2a + 2b + 2c + 2d + 2e$$

$$e = b$$




Formulas for perimeters (II)

Two squares and one rectangle in between:



- Give formulas for the perimeters P , Q and R related to a and/or b :
 $P = \dots\dots\dots$ $Q = \dots\dots\dots$ $R = \dots\dots\dots$
- Explain the relationship: $Q = \frac{1}{2}P + \frac{1}{2}R$

Three hexagons with only angles of 120° :



- Give formulas for P , Q and R :
 $P = \dots\dots\dots$ $Q = \dots\dots\dots$ $R = \dots\dots\dots$
- Explain why $Q = \frac{2}{3}P + \frac{1}{3}R$ is valid.
- Design a hexagon with angles of 120° and perimeter S in such a way that the following relationship is valid: $S = \frac{1}{3}P + \frac{2}{3}R$

Polynomials and weights (I)

$$a + a + b + b + b + c + c + c + c = 2a + 3b + 4c$$

nine terms

trinomial


2, 3 and 4 are the **weights** of a , b and c

With the weights 2, 3 and 4 and the letter a , b and c you can make other trinomials like for example: $4a + 3b + 2c$.

There are totally six different trinomials in a , b , c with weights 2, 3, 4.

- Write down the other four trinomials.
- Add the six trinomials. Which trinomial is the result?

If it is known that $c = b$, you can make a binomial of $2a + 3b + 4c$.



You can do the same with the other five trinomials.

- How many different binomials do you get? Which ones?
- If you also know that $a = b$ you can simplify the binomials further. What is the result?

Polynomials and weights (II)

$$w + w + x + y + y + y + y + z + z = 2w + x + 5y + 2z$$

quadrinomial

w has a weight 2, x a weight 1, y a weight 5 and z a weight 2

The weight 1 is often left out in polynomials but if you think it's more clear, you may write: $2w + 1x + 5y + 2z$


The sum of the weights in this polynomial is 10.

- Give five other quadrinomials in w , x , y and z , for which the sum of the weights is equal to 10.

- Add those five quadrinomials.

What is the sum of the weights of the resulting quadrinomial?

If it is known that $w = x$ and $z = y$, then $2w + x + 5y + z$ can be simplified to a binomial:



Simplify your quadrinomials to binomials, when given: $w = x$ en $z = y$.

- Which binomials do you get?

- The sum of all these binomials is equal to a binomial. Which one?

Polynomials and weights (III)

In a college are held three tests for math in the last period before summer. Two tests are held in the 'normal' time (1 hour) and for one test the students have 2 hours).

To determinate the final score it's not fair to give the three results the same weight.

The teacher tells the students how he will calculate the final score:

$$\frac{A + B + 2C}{4}$$

In this formula A is the score for the first, B for the second one and C for the third (long) test.

Another formula which gives the same result, is:

$$\frac{1}{4}A + \frac{1}{4}B + \frac{1}{2}C$$

Tanja has for the first test the score 4, for the second one the score 6 and for the last one the score 9.

- Calculate her final score, using both formulas.
- Calculate the final result for some other scores.

The teacher calculated a so called **weighted mean**.

Another weighted mean of A , B en C is for instance:

$$\frac{A + 2B + 3C}{6} \quad \text{or:} \quad \frac{1}{6}A + \frac{1}{3}B + \frac{1}{2}C$$

- Suppose that a teacher uses this formula, to determinate the final score of three tests. Can this formula be fair?
- Give two formulas for the 'normal mean' (each test has the same 'weight').


Equivalent (I)

$\frac{A + B + 2C}{4}$ and $\frac{1}{4}A + \frac{1}{4}B + \frac{1}{2}C$ are equivalent

That means:

whatever the numbers are that you substitute for A , B and C , the resulting values of both expressions will be the same.

This equivalency can be understood looking at this diagram:



● Equivalent or not?

$$\frac{A + 2B + 3C}{6} \stackrel{?}{=} \frac{1}{6}A + \frac{1}{3}B + \frac{1}{2}C$$

$$\frac{A + 2B + 3C}{6} \stackrel{?}{=} \frac{A}{6} + \frac{B}{3} + \frac{C}{2}$$

$$2 \times (5A + 3B + C) \stackrel{?}{=} 10A + 3B + C$$

$$2 \times (5A + 3B + C) \stackrel{?}{=} 2 \times (5A + 3B) + C$$

$$2 \times (5A + 3B) + C \stackrel{?}{=} 10A + 6B + C$$

Equivalent (II)

Here are ten expressions

A group of *equivalent* expressions do we call a *family*.

- Connect the members of the same family by a line.

$3X + 18Y + 63Z$

$3 \times (X + 6Y + 21Z)$

$3 \times (X + 7Z) + 6Y$

$3X + 3 \times (2Y + 7Z)$

$3X + 18Y + 21Z$

$3 \times (X + 6Y) + 21Z$

$3X + 6Y + 21Z$

$3 \times (X + 2Y + 7Z)$

$3 \times (X + Y) + 3 \times (Y + 7Z)$

Equivalent (III)

- Find as many expressions as you can which are equivalent with:


$$**5a + 10b + 20c**$$

Rule: each of the letters a , b and c may appear only once in each expression.

- The same for:

$$\frac{P + 2Q + 2R + 4S}{12}$$


Red-white-blue (I)



The number strip has a repeating pattern red-white-blue-red-white-blue, etc.


- ◆ What colour has the cell of number **100**? The same question for **1000**?
- ◆ Which of the numbers between **1000** and **1010** are in a red cell?
- ◆ Find a 'blue number' of five digits.
- ◆ If you add two 'red numbers' , you always get a red number.
Do you think this true? Why??
- ◆ What can you say about the sum of two 'blue numbers' ?

Red-white-blue (II)




Below you see separate strips of the red, white and blue numbers .


◆ Give an expression to each number strip.



Red-white-blue (III)



The red, white and blue numbers can be represented by dot patterns.
For example:



From the dot patterns you can understand: **White + Blue = Red**

◆ Explain this.

◆ Complete the chart for all combinations of colours.

+	Red	White	Blue
Red			
White			Red
Blue			

Different differences (I)

Isabelle did a job and earned € 100. She wants to buy a pair of sport shoes. They normally cost € 70 . So she expects € 30 will be left.

How lucky she is! Entering the shop she discovers that the pair she wants is reduced by € 8.

- How many Euro's does she have left?
- There are two ways to calculate this:

a) $100 - (70 - 8)$ and b) $(100 - 70) + 8$

Which method a) or b) did you use?

Explain, without looking at the result, that the alternative method is just as well.

Suppose that Isabelle already knew, that the price of the shoes was reduced, but she did not know how much ...

So she knew that more than € 30 should be left!

- Use the story to explain: $100 - (70 - a) = 30 + a$

A more general equality: $100 - (p - a) = 100 - p + a$

- Explain this equality.
(You may suppose $p < 100$ and $a < p$).

A frequently occurring error: $100 - (p - a) = 100 - p - a$

- Invent a short story in which someone has $100 - p - a$ left from 100 Euros in stead of $100 - p$.

- Fill in the right expression: $100 - (\dots\dots\dots) = 100 - p - a$

Different differences (II)

If you subtract **less**, there will be left **more**.

Example:

$$\begin{array}{l} 50 - 20 = 30 \\ 50 - (20 - x) = 30 + x \end{array}$$

If you subtract **more**, there will be left **less**.

- Invent an example:

- Four conclusions. The first one is complete. Explain those.
- Complete the other three.

$$\begin{array}{l} y = 10 - x \\ z = 15 - y \end{array}$$

$$z = 5 + x$$

$$\begin{array}{l} y = 10 - x \\ z = 15 + y \end{array}$$

$$z = \dots\dots\dots$$

$$\begin{array}{l} y = 10 + x \\ z = 15 + y \end{array}$$


$$z = \dots\dots\dots$$

$$\begin{array}{l} y = 10 + x \\ z = 15 - y \end{array}$$


$$z = \dots\dots\dots$$

- Invent some other conclusions in the same style..
Use other symbols than x, y, z .

Different differences (III)



- The sum of INPUT and OUTPUT is 100 in each case. Check this!
- Complete the chain :



The chain on the left corresponds to the following chain of differences:

$$30 - [80 - (150 - (100 - x))] = \dots$$

- This expression is equivalent with a very simple one. Which one?
- Make chains corresponding to:

$$10 - (9 - (8 - y))$$

$$16 - [9 - (4 - (1 - a))]$$

$$32 - [16 - [8 - (4 - (2 - k))]]$$
- What are the three resulting binomials?
- Invent such a 'chain-exercise' by yourself.

Let it be true (I)

$$\begin{array}{c} X = 5 \\ \downarrow \\ 20 + 5X = 45 \end{array}$$

Check that this is true.

- Find a value for X that makes this true:

$$\begin{array}{c} X = \dots \\ \downarrow \\ 20 + 5X = 35 \end{array}$$

- Find in each of the following cases a value for X that makes the conclusion true.
You may 'guess smartly' and check by calculating!

$$\begin{array}{c} X = \dots \\ \downarrow \\ 25 + X = 125 \end{array}$$

$$\begin{array}{c} X = \dots \\ \downarrow \\ 25X = 125 \end{array}$$


$$\begin{array}{c} X = \dots \\ \downarrow \\ \frac{1}{X+1} = \frac{1}{5} \end{array}$$

$$\begin{array}{c} X = \dots \\ \downarrow \\ \frac{1}{X-1} = \frac{1}{5} \end{array}$$


$$\begin{array}{c} X = \dots \\ \downarrow \\ X \times (X + 1) = 30 \end{array}$$

$$\begin{array}{c} X = \dots \\ \downarrow \\ \frac{X}{X+1} = \frac{3}{4} \end{array}$$

Let it be true (II)



Let it be true (III)



- For which values of ***b, c, d, e, f, g*** all equalities in the triangle are valid?
- How do the answers change if ***a*** is not equal to 4, but to 8?

Generation problems

Today, Anja, her mother, her grandmother and her great-grandmother are together 200 years old.

Anja's mother was 30 years old when Anja was born.

Grandma was 25 years old when Anja's mother was born.

Great-grandma was 20 years old at the birth of Anja's grandma.

Suppose the age of Anja, her mother, her grandmother and her great-grandmother respectively equal to ***a***, ***b***, ***c*** and ***d*** years.

- Write down everything you know about ***a***, ***b***, ***c*** and ***d*** in formulas.

- Calculate ***a***, ***b***, ***c*** and ***d***.

Peter, his father, his grandfather and his great-grandfather also are together 200 years old.


Father is 4 times as old as Peter, grandfather is $1\frac{1}{2}$ times as old as father, great-grandfather is $1\frac{1}{2}$ times as old as grandfather.

Suppose the age of Peter, his father, his grandfather and his great-grandfather are respectively equal to ***p***, ***q***, ***r*** and ***s*** years.

- Write down everything you know about ***p***, ***q***, ***r*** and ***s*** in formulas.

- Calculate ***p***, ***q***, ***r*** and ***s***.

On the number line (I)



Between 1994 and 2003 are 9 years.


- How many years are there between 2011 and 1945?

In the year n astronauts from Earth land on Mars for the first time.

One year later they return to Earth. That will be in the year $n + 1$.

Again one year later the astronauts take an exhibition about their trip, around the world. That will be in the year $n + 2$.

The construction of the launching rocket began one year before the, landing on Mars, so this was in the year $n - 1$



Between $n - 1$ and $n + 2$ there are 3 years.

You may write:

$$(n + 2) - (n - 1) = 3$$

- How many years are there between $n - 4$ and $n + 10$?

- Calculate:

$$(n + 8) - (n - 2) =$$

$$(n + 7) - (n - 3) =$$


$$(n - 1) - (n - 4) =$$

$$(n + 3) - (n - 3) =$$

- How many years are there between $n - k$ and $n + k$?

On the number line (II)


Even and odd years



An **arbitrary** even year can be represented by $2n$.


In two years it will be the year $2n + 2$, that is the even year that follows the even year $2n$.

The even year that comes before $2n$ is the year $2n - 2$.



- What is the even year that follows the year $2n + 2$? And what is the even year that comes before the year $2n - 2$?

The odd years are between the even years.




- Write expressions for the odd years on the number line.
- Calculate:

$$(2n + 8) - (2n - 6) =$$

$$(2n + 3) - (2n - 3) =$$

$$(2n + 4) - (2n - 3) =$$

Olympic years are divisible by 4.




- How can you represent an **arbitrary** Olympic year?
- Which Olympic year succeeds that year and which precedes it?

Olympic wintergames are presently held in a year that exactly lies between two Olympic years.


- How can you represent an **arbitrary** year of wintergames?

On the number line (III)




The number x is chosen in such a way that $x - 8 = 20 - x$ is valid.

- Which value has x ?
- Which value has x if $x - 1971 = 2001 - x$?




● $x - n = 3n - x \longrightarrow x = \dots\dots\dots$ expression in n

● $x - 5n = 25n - x \longrightarrow x = \dots\dots\dots$ expression in n



● $x - n = m - x \longrightarrow x = \dots\dots\dots$ expression in n and m


On the number line (IV)



The number y is chosen in such a way that: $y - 8 = 2 \times (20 - y)$

● Which value has y ?

● Which value has y if: $y - 1971 = 2 \times (2001 - y)$?



$y - n = 2 \times (m - y)$

● Which expression (in n and m) can you find for y ?


Dot patterns (I)

Nicomachos lived about the year 100 AC in Greece.

He wrote a book about the 'admirable and divine properties of natural numbers.


Nicomachos sometimes used dot patterns to represent numbers.

Below you see the most famous examples:



He gave every type a geometrical name. .


To begin with, consider the family of the *square numbers*.



- Write the next ten square numbers. You need not to draw the corresponding patterns (but you can 'see' them in your mind).
- Consider the steps between consecutive square numbers. Do you see any rule? How can you see that rule in the dot patterns?
- 144 is a square number. How about 1444? And 14444? Use a calculator to investigate this.

Dot patterns (II)

The first four *oblong numbers*



- Write the next ten oblong numbers. Continue the dot patterns in your mind.
- Consider the steps between successive square numbers.. Do you see any rule? How can you see that rule in the dot patterns?
- Is 9900 an oblong number? Explain your answer.

Take the mean of pairs of consecutive oblong numbers.

So:

the mean of 2 and 6 is 4
the mean of 6 and 12 is 9
the mean of 12 and 20 is 16


- Continue this at least five times.
Which special numbers do you get as result?
Try to explain your discovery.

Dot patterns (III)

Nicomachos was not the first person who used dot patterns for numbers. 600 years before (about 500 BC) lived Pythagoras, a scholar who was the leader of a religious sect.

In the doctrine of Pythagoras 'natural numbers' played the leading part. The device of him and his disciples was: '*everything is number*'.


Their favorite number was 10, being the sum of 1, 2, 3 and 4.



10 is the fourth number in the sequence of the *triangular numbers*.

- Write the next ten triangular numbers.


The dot patterns of the triangular numbers can also be drawn in this way:




- Which special numbers do you get if each of the triangular numbers are doubled? How can you explain this using the dot patterns?
- Is 4950 a triangular number? Explain your answer.
- Calculate the sum of all natural numbers smaller than 100.

Strips and dots (I)

square numbers



oblong numbers



The expression for the square numbers is mostly written as: n^2

Pronounce: n square.


- Compare both strips. Which strip can you add to the left one to get the right one?

The expressions $n \times (n + 1)$ and $n^2 + n$ are equivalent.


- How can you explain this using dot patterns?

Strips and dots (II)

triangular numbers



oblong numbers



Compare the triangular numbers with the oblong numbers.


- Whic expression fits the strip of triangular numbers?
- Give one (or more) expressions which are equivalent with this.

Using the expression for triangular numbers you can calculate the sum of the first hundred natural numbers:


- $1 + 2 + 3 + 4 + 5 + \dots + 98 + 99 + 100 =$ _____

Strips and dots (III)

triangular numbers



pentagonal numbers




Compare the numbers of both strips.

- Which pentagonal number succeeds 92?
- Find an expression which represents the sequence of pentagonal numbers.

Number spiral (I)

You can make a number line in the shape of a spiral!



The black vertices of the spiral correspond with special families of numbers.

- Which families?


You can see in the diagram that every square number lies exactly in the middle between two oblong numbers.

For example: **49** lies in the middle between **42** and **56**.

Of course, for **49 = 7 × 7** and **42 = 6 × 7** and **56 = 8 × 7**

- The square number 144 lies in the middle between the oblong numbers and
- The square number 1444 lies in the middle between the oblong numbers and
- The square number n^2 lies in the middle between the oblong numbers and

Number spiral (II)



Now the dots on the vertical parts of the number spiral are black, the others white. If you add the black numbers form one vertical part, the result will be equal to the sum of the succeeding white numbers.

You can check that:

$$1 + 2 = 3$$

$$4 + 5 + 6 = 7 + 8$$

$$9 + 10 + 11 + 12 = 13 + 14 + 15$$


$$16 + 17 + 18 + 19 + 20 = 21 + 22 + 23 + 24$$

- What will be the next line of this scheme?
- You can check this line without calculating both sums.
Hint: Mark the steps from 'black' to 'white'.

Line n begins with the black number n^2 .


- Give an expression for the last black number on that line? How big are the he steps from 'black' to 'white' .?
- Try to explain: 'black sum' = 'white sum'

Geometric Algebra (I)




$$h \times a + h \times b = h \times (a + b)$$
 or

$$ha + hb = h(a + b)$$



$$10a + 10b = \dots\dots\dots$$

.....











.....


.....








Geometric Algebra (II)



Geometric Algebra (III)









$pq - 2rs$



- Invent a figure with area $ab - 3c^2$
- Also one with area $p^2 + 4q^2$

Geometric Algebra (IV)



Isn't it remarkable? (I)

a and b are positive integers with the sum 10

S is the sum of the squares of a and b , so: $S = a^2 + b^2$


T is the sum of the ab and ba , so: $T = ab + ba$

- Which values can S have? And T ? Fill in the table.

$a + b = 10$		S	T
a	b	$a^2 + b^2$	$ab + ba$


- Calculate the sum $S + T$ in each case. What do you discover?
- Investigate what happens with $S + T$ if a and b are decimal numbers with sum 10, for instance $a = 3,8$ and $b = 6,2$. Investigate some other examples. What did you find?

How remarkable? (II)



- Draw a square with side a . Also one with side b .
- Draw a rectangle with horizontal side a and vertical side b . Also one with horizontal side b and vertical side a .
- How can you explain that $(a^2 + b^2) + (ab + ba) = 100$?

Strips and expressions (I)



- Fill in the missing numbers on the strips.
- Equivalent or not?

$$\boxed{n^2} \stackrel{?}{=} \boxed{n \times n}$$

$$\boxed{2n + 1} \stackrel{?}{=} \boxed{n \times n + 1}$$

$$\boxed{n^2 + 1} \stackrel{?}{=} \boxed{n^2 + 1^2}$$

$$\boxed{2(n + 1)} \stackrel{?}{=} \boxed{2n + 1}$$



$$\boxed{(n + 1)^2} \stackrel{?}{=} \boxed{n^2 + 1^2}$$

$$\boxed{(n + 1)^2} \stackrel{?}{=} \boxed{n^2 + 2n + 1}$$



$$\boxed{n^2 + 1} \stackrel{?}{=} \boxed{n \times n + 1 \times 1}$$

$$\boxed{(n + 1)^2} \stackrel{?}{=} \boxed{n^2 + 1 + 2n}$$


Strips and expressions (II)




Formulas about squares



$$\longrightarrow (n + 1)^2 = n^2 + 2n + 1$$



$$\longrightarrow (n + 2)^2 = n^2 + 4n + 4$$




$$\longrightarrow (n + 3)^2 = n^2 + 6n + 9$$

- How will this continue? Write the next three formulas..

Powerful tables (I)

Below you see a table with powers of 2:

n	2^n
0	1
1	2
2	4
3	8
4	16
5	32
6	64
7	128
8	256
9	512
10	1024
11	2048
12	4096
13	8192
14	16384
15	32768
16	65536
17	131072
18	262144
19	524288
20	1048576



- Find the results of the following products using the table:
 $16 \times 8192 = \dots$
 $8 \times 16384 = \dots$
 $512 \times 512 = \dots$
 $1024 \times 1024 = \dots$
- Write all pairs of two positive integers with a product equal to
 1048576

Powerful tables (II)

Table with powers of 3	
n	3^n
0	1
1	3
2	9
3	27
4	81
5	243
6	729
7	2187
8	6561
9	19683
10	59049
11	177147
12	531441
13	1594323
14	4782969
15	14348907
16	43046721
17	129140163
18	387420489
19	1162261467
20	3486784401

fill in:

$$243 \times 81 = \dots\dots\dots$$

$$\begin{array}{ccc} \downarrow & & \downarrow \\ 3^{\dots} & \times & 3^{\dots} \\ \uparrow & & \uparrow \\ 3^{\dots} & \times & 3^{\dots} = 3^{\dots} \end{array}$$

- Find the results of the following products using the table:

$$81 \times 19683 = \dots$$

$$2187 \times 59049 = \dots$$

$$6561 \times 6561 = \dots$$

$$729 \times 729 \times 729 = \dots$$

- Find the results of the following powers using the table:

$$81^3 = \dots$$

$$243^4 = \dots$$

$$27^5 = \dots$$

- Which number is smaller: 9^{10} or 10^9 ?

Powerful tables (III)

- Make a table with powers of 5 (until 5^{10})
Design some problems which you can solve using this table.

n	5^n
0	1
1	5
2	
3	
4	
5	
6	
7	
8	
9	
10	


- The table with powers of 1 is very simple. Why?
- Do you know a table of powers with sharp rising results, which is very easy to write? Which one?
- If you put the tables with powers of 2 and 3 next to each other, and if you multiply the numbers on the same line, you get:

$$\begin{aligned}1 \times 1 &= 1 \\2 \times 3 &= 6 \\4 \times 9 &= 36 \\8 \times 27 &= 216 \\&\text{etc.}\end{aligned}$$

The results are just the powers of 6.
You can check this using your calculator.

- How can you explain without calculator: $2^{10} \times 3^{10} = 6^{10}$?

Divisors(I)



A slab of chocolate contains 12 pieces.

- For what numbers of persons fair sharing in whole pieces is possible?

If you gave a correct answer to the last question, then you found the divisors of 12. Maybe you forgot that it's possible to eat the slab alone, but also 1 is a divisor of 12.

- Complete the following divisor-table:

number	divisors	number of divisors
1	1	1
2	1 , 2	2
3	1 , 3	2
4	1 , 2 , 4	3
5	1 , 5	2
6		
7		
8		
9		
10		
11		
12	1 , 2 , 3 , 4 , 6 , 12	6
13		
14		
15		
16		
17		
18		
19		
20		
21		
22		
23		
24		
25		

Divisors (II)

Consider the divisor-table of the numbers 1, 2, 3, ..., 25.

There are numbers with only two divisors

Such numbers are called **prime numbers**.

- What are the prime numbers below 25?
- What are the next three prime numbers?

Maybe you discovered already a smart way to find the divisors of a number. If you know one divisor, you can (mostly) find a second one immediately. So you can find pairs of divisors.

Take for example the number 108.


The first pair of divisors is trivial: 1 and 108.

Because 2 is a divisor, you have immediately 54 (for $2 \times 54 = 108$).

The next divisor is 3, and you have also 36 (for $3 \times 36 = 108$).

Etcetera.

So we have: 1, 2, 3, 4, 6, 9, **12**, **18**, **27**, **36**, **54**, **108**



The bold numbers do you get as a present more or less.

- In the case of 108, you only have to 'try' the numbers 1 to 10. Explain this..
- Find in this way the divisors of: 88 ; 144 ; 210

The majority of numbers have an even number of divisors.

- For which type of numbers, the number of divisors is odd?

In the table with powers of 2 you can find the 8192.

- Which divisors has this number?
- How many divisors has 3^{10} ?
- Find a big number by yourself, of which you quickly can give the number of divisors'.

Divisors (III)


Of two natural numbers you know:

* **their product is 80**


* **their sum is 21**

- Which are the two numbers?

- Find the two natural numbers



- Find the two natural numbers



- Find a value for n in such a way that $n \times (n - 4) = 77$

- Find a value for p in such a way that $(p + 7) \times (p + 10) = 108$

Prime numbers

The Greek scholar Eratosthenes lived about 240 before Chr.
Eratosthenes was a universal man ; besides a mathematician he also was, historian, geographer, philologist and poet. He is famed for his measurement of the earth.
Furthermore he found a smart way to find prime numbers

Look at the 'hundred-chart' below.

1 is no prime number, so the cell of 1 is shaded.

2 is a prime number , the cell of 2 remains white.

Then all the cells of multiples of 2 are shaded.

1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20
21	22	23	24	25	26	27	28	29	30
31	32	33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48	49	50
51	52	53	54	55	56	57	58	59	60
61	62	63	64	65	66	67	68	69	70
71	72	73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88	89	90
91	92	93	94	95	96	97	98	99	100

3 is a prime number; the cell of 3 remains white.

* Now shade the cells of all multiples of 3, which are not already shaded.

5 is a prime number; the cell of 5 remains white.

* Now shade the cells of all multiples of 5, which are not already shaded.

7 is a prime number; the cell of 7 remains white.

* Now shade the cells of all multiples of 7, which are not already shaded

Now all white cells are containing a prime number!!

* How can you explain this?

Hint: consider the prime number after 7, that is 11. What are the multiples of 11 smaller than 100? Why can you be sure without looking at the chart that the cells of these numbers are already shaded?

Prime factors


Prime numbers can be considered as the atoms for the natural numbers.

Every natural number is equal to a product of prime numbers.

For example: $540 = 2 \times 2 \times 3 \times 3 \times 5$ or shortened: $540 = 2^2 \times 3^3 \times 5$

We say: 540 is *decomposed in factors* or *factorized*.

You can find such a decomposition in many ways. Try to split the number in two factors (bigger than 1 and smaller than the given number); if possible you do the same with both factors, and you repeat this process until you only have prime factors. For example with 540:



There are many other 'trees of decomposition'; the final result of all these trees has to be the same!

- Factorize in this way the following numbers:

135

135 =

704

704 =

2100

2100 =

- Choose three other numbers by yourself and find the decomposition in prime factors.

Amicable numbers (I)


To clinch a close friendship between two persons, sometimes both friends wear an amulet or a ring with a short text inscribed on it.

In the age of Pythagoras (500 BC) they used sometimes the numbers **220** and **284** for such an inscription.

These two numbers are called **amicable numbers**.

The mystery of the numbers 220 and 284 has something to do with their divisors.

- Find all divisors of 220.
Add up all these numbers except 220..
- Do the same with the divisors of 284 (except 284).



If you did not make a mistake, you found a remarkable fact:

the sum of the divisors (except 220) of 220 = 284
the sum of the divisors (except 284) of 284 = 220

This property of a pair of numbers is very exceptional.

In later times famous mathematicians made a great effort to find other pairs of amicable numbers.

The French mathematician Pierre Fermat, who actually was a lawyer, discovered in 1636 a new pair: **17296** en **18416**.

Leonard Euler, a Swiss mathematician, whose portrait is on the Swiss banknotes, published in 1747 a list with thirty pairs of (big) amicable numbers; a few years later this list was extended to sixty pairs.

It's remarkable that all famous scholars didn't discover a rather small pair, namely **1184** and **1210**.

This pair was found by sixteen years old Italian boy, named Nicolo Paganini, in the year 1866.

- Check that **1184** and **1210** are amicable numbers.

Amicable numbers (II)

The Arabic mathematician Tabit ibn Qorra, who lived from 826 to 901, found a recipe to make pairs of amicable numbers.

His recipe was rather complicated as you can see below.

IF *and* *IF*

$$p = 3 \times 2^n - 1$$

$$q = 3 \times 2^{n-1} - 1$$

$$r = 9 \times 2^{2n-1} - 1$$

p, q, r
are three odd
prime numbers

THEN $A = 2^n \times p \times q$

 $B = 2^n \times r$
are AMICABLE numbers

- Take $n = 2$. Check if p, q, r indeed are prime numbers.
Use the values of p, q and r to calculate A and B .
Now you see ...
- Take $n = 3$. Which of the numbers p, q, r is **not** a prime number.

For $n = 3$ the recipe of Tabit ibn Qorra don't give amicable numbers!
For $n = 4$ it does! If you don't have a long list of prime numbers, it is difficult to check if p, q, r are prime in this case. A computer has validated that this is true.

- Calculate A and B for $n = 4$ and compare the result with the discovery of Pierre de Fermat (previous page).
- The numbers of Nicolo Paganini can not be made by the recipe above.
Factorize these numbers and explain why not.

Obviously the formula of Tabit ibn Qorra don't give all pairs of amicable numbers !

Operating with powers (I)

$$a \times a \times a \times b \times b \times c = a^3 \times b^2 \times c^1 = a^3 b^2 c$$


The **exponents** of a , b , and c are 3, 2 and 1.
(note: the exponent 1 is mostly not written)

With the exponents 3, 2 and 1 and the letters a , b and c also can be made other products, for example: ab^3c^2 .

There are six different products that one can make using a , b , c and the exponents 1, 2, 3.

- Write the other four products.
- Multiply the six products to each other.
The result can be written in the form $a \cdots b \cdots c \cdots$
Which exponents do you get?

If it is known that $b = a$, you can simplify $a^3 b^2 c$



You can do the same with the other five products.

- How many different products do you get? Which ones?
- If you also know that $c = a$ you can write each of these products as a **power** of a . Which one?

Operating with powers (II)

$$2m^3 \times 3m^2 = 2 \times 3 \times m^5 = 6m^5$$

$$2 \times m \times m \times m$$

$$3 \times m \times m$$

- Find other multiplications, as many as possible, with the same result:

$$\dots \times \dots = 6m^5$$

$$\dots \times \dots = 6m^5$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Operating with powers (III)

$$3a = a + a + a$$

$$a^3 = a \times a \times a$$

$$\leftarrow a = z^2 \rightarrow$$

$$3z^2 = z^2 + z^2 + z^2$$

$$(z^2)^3 = z^2 \times z^2 \times z^2 = z^6$$

$$\boxed{4z^2} + \boxed{3z^2} + \boxed{2z^2} - \boxed{z^2} = \boxed{8z^2}$$

- On the places _____ you can fill in + , - or \times to make the following equalities:

$$\boxed{4z^2} \text{ — } \boxed{3z^2} \text{ — } \boxed{2z^2} \text{ — } \boxed{z^2} = \boxed{10z^2}$$

$$\boxed{4z^2} \text{ — } \boxed{3z^2} \text{ — } \boxed{2z^2} \text{ — } \boxed{z^2} = \boxed{24z^8}$$


$$\boxed{4z^2} \text{ — } \boxed{3z^2} \text{ — } \boxed{2z^2} \text{ — } \boxed{z^2} = \boxed{14z^4}$$

$$\boxed{4z^2} \text{ — } \boxed{3z^2} \text{ — } \boxed{2z^2} \text{ — } \boxed{z^2} = \boxed{10z^4}$$

$$\boxed{4z^2} \text{ — } \boxed{3z^2} \text{ — } \boxed{2z^2} \text{ — } \boxed{z^2} = \boxed{2z^2}$$

Between two squares

A rectangle between two squares:



- Fill in expressions in a and b : $P = \dots\dots\dots$, $Q = \dots\dots\dots$, $R = \dots\dots\dots$

Below you see six formulas with P , Q and R .

- Investigate which formulas are correct.

$$Q^2 = PR$$

$$Q = \frac{P+R}{2}$$


$$Q = \sqrt{PR}$$

$$2Q = P+R$$

$$Q - P = R - Q$$

$$\frac{Q}{P} = \frac{R}{Q}$$

Between two cubes



- Fill in expressions in a and b : $P = \dots\dots\dots$, $Q = \dots\dots\dots$, $R = \dots\dots\dots$, $S = \dots\dots\dots$
- Check the following formulas:


$$Q^2 = PR$$

$$Q = \sqrt{PR}$$

$$\frac{Q}{P} = \frac{R}{Q}$$

- Invent at least three formulas with only Q , R and S .
- Also invent some formulas with P , Q , R and S .

You can count on it (I)




- Check the calculations on the blackboard. Continue the sequence with some more lines. What do you think?
- Give some other calculations, fitting in the sequence, with numbers between 100 and 1000. Check if the result will be 2?


This is the general rule:

$$(n + 1) \times (n + 2) - n \times (n + 3) = 2$$

- How can you explain the validity, using the pictures in the cloud?




You can count on it (II)




- What is the regularity in this sequence of calculations?
- Which formula corresponds with this sequence?

- Draw a picture that explains the formula:



You can count on it (III)




- Design a similar sequence of calculations (with the same result on each line).


- Geef daar ook de passende formules bij.

You can count on it (IV)

$$(n + 1) \times (n + 2) = n^2 + 3n + 2$$



$$n \times (n + 3) = n^2 + 3n$$



$(n + 2) \times (n + 3) = \dots\dots\dots$

$n \times (n + 5) = \dots\dots\dots$

-

$(n + 2) \times (n + \dots) = \dots\dots\dots$

$n \times (n + \dots) = \dots\dots\dots$

- 10

$\dots\dots\dots$

$\dots\dots\dots$

-

Dictation

in words

in symbols

the sum of n and m is reduced with 5

$$n + m - 5$$

the sum of n and 8 is multiplied by n

$$(n + 8) \times n$$

the product of n and m is reduced with k

the sum of the squares of n and m

the square of the sum of n and m

the sum of 2 times n and 3 times k

the sum of n and 6 is multiplied by the difference of n and 6

the product of the third powers of n and m

the square of 5 times n

5 times the square of n

The price of algebra (I)

Algebra takes time, and time is money.
Below you see a detailed 'price' list

Price list:	
operations +, -, ×, :, /	1 point each time
taking a square	2 points each time
taking the 3rd power	3 points each time
taking the 4th power	4 points each time
etc.	etc.
using variables	1 point each time
parenthesis and numbers	free

Example 1: what is the price of $3n + m$?

3	number	free
n	using variable	1 point
$3 \times n$	multiplication	1 point
m	variable	1 point
$3 \times n + m$	addition	1 point
total price		4 points

Example 2: what is the price of $(3n + m)^2$?

$3n + m$	just calculated	4 points
$(3n + m)^2$	square	2 points
total price		6 points

- Find the price of:

$$n^2 + 3n$$

$$n \times (n + 3)$$

$$(n + 1) \times (n + 3)$$

$$n^2 + 4n + 3$$

The price of algebra (II)

$n^2 + 3n$ and $n \times (n + 3)$ are equivalent expressions.

They have not the same prices yet! (See preceding page).

- Here are pairs of equivalent expressions. Make sure of it. Find out for each pair which of the two is the cheapest..

$$n + n + n + n \quad \text{and} \quad 4 \times n$$

$$n \times n \times n \times n \quad \text{and} \quad n^4$$

$$n + n + n + n \quad \text{and} \quad 2 \times n + 2 \times n$$

$$(m + 1)^2 \quad \text{and} \quad (m + 1) \times (m + 1)$$

$$(m + 1)^2 \quad \text{and} \quad m^2 + 2m + 1$$

$$a^2 \times a^3 \quad \text{and} \quad a^5$$

$$(a^3)^2 \quad \text{and} \quad a^6$$

The price of algebra (III)

- Compare the prices of a^4b^4 and $(a^2b^2)^2$

Both expressions are equivalent, but the second one is 2 points cheaper.

- Try to find an expression as cheap as possible, which is equivalent with a^4b^4

You can write n^{15} in various ways (it's to say: replacing by an equivalent expression).

Here are some possibilities:

$$n \times n \times n \times n \times n \times n \times n \times n \times n \times n \times n \times n \times n \times n \times n$$

$$(n \times n \times n \times n \times n)^3$$

$$n^{10} \times n^5$$

- Which of them is cheaper than n^{15} ?
Try to find an equivalent expression with the lowest price.

- Find the cheapest expression equivalent with x^{24}

Splitting fractions (I)

Unit fractions

4000 years ago the mathematicians in Egypt worked with unit fractions, like:

$$\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5} \text{ etc.}$$

Thus with fractions with a **numerator** equal to 1.

A fraction with another numerator than 1, always can be split up in unit fractions. This one is easy:

$$\frac{3}{4} = \frac{1}{4} + \frac{1}{4} + \frac{1}{4}$$

It is less insipid if it is demanded to use as few unit fractions as possible, in this case:

$$\frac{3}{4} = \frac{2+1}{4} = \frac{1}{2} + \frac{1}{4}$$

Another example:

$$\frac{19}{24} = \frac{12+6+1}{24} = \frac{1}{2} + \frac{1}{4} + \frac{1}{24}$$

- Try to split into unit fractions, as few as possible:

$$\frac{7}{8} = \frac{\dots + \dots + \dots}{8} = \frac{1}{\dots} + \frac{1}{\dots} + \frac{1}{\dots}$$

$$\frac{5}{6} = \dots = \dots$$

$$\frac{13}{16} = \dots = \dots$$

$$\frac{13}{18} = \dots = \dots$$

$$\frac{99}{100} = \dots = \dots$$

Splitting fractions (II)

Numerator and denominator of a fraction can be divided (or multiplied) by the same number; the value of the fraction does not change then.

Examples: $\frac{5}{15} = \frac{1}{3}$ (numerator and denominator are divided by 3)

$$\frac{b}{5b} = \frac{1}{5} \quad (\text{numerator and denominator are divided by } b)$$

This rule is applied with splitting into unit fractions.

Examples: $\frac{n+1}{3n} = \frac{n}{3n} + \frac{1}{3n} = \frac{1}{3} + \frac{1}{3n}$

$$\frac{a+b}{ab} = \frac{a}{ab} + \frac{b}{ab} = \frac{1}{b} + \frac{1}{a}$$

- Split into unit fractions as few as possible :

$$\frac{x+y}{xy} =$$

$$\frac{k+m+n}{kmn} =$$

$$\frac{p+1}{pq} =$$



$$\frac{p+1}{p^2} =$$

- Fill in the correct expressions:

$$\frac{\dots + \dots + \dots + \dots}{abcd} = \frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d}$$

Fractions on strips (I)

Fill the empty cells:



Fractions on strips (II)

Fill the empty cells:


1	\times	$\frac{1}{2}$	$=$	$\frac{1}{2}$
$\frac{1}{2}$		$\frac{1}{3}$		$\frac{1}{6}$
$\frac{1}{3}$		$\frac{1}{4}$		$\frac{1}{12}$
$\frac{1}{4}$		$\frac{1}{5}$		$\frac{1}{20}$
$\frac{1}{5}$		$\frac{1}{6}$		$\frac{1}{30}$

$\frac{1}{n}$	\times	$\frac{1}{n+1}$	$=$	
---------------	----------	-----------------	-----	--

1	$:$	$\frac{1}{2}$	$=$	2
$\frac{1}{2}$		$\frac{1}{3}$		$\frac{3}{2}$
$\frac{1}{3}$		$\frac{1}{4}$		$\frac{4}{3}$
$\frac{1}{4}$		$\frac{1}{5}$		$\frac{5}{4}$
$\frac{1}{5}$		$\frac{1}{6}$		$\frac{6}{5}$



$\frac{1}{n}$	$:$	$\frac{1}{n+1}$	$=$	
---------------	-----	-----------------	-----	--

Multiplying fractions (I)



- Draw a picture to show: $\frac{4}{5} \times \frac{2}{3} = \frac{4 \times 2}{5 \times 3} = \frac{8}{15}$
- Calculate: $\frac{3}{5} \times \frac{1}{4} =$ $\frac{3}{5} \times \frac{4}{5} =$
 $\frac{3}{4} \times \frac{1}{5} =$ $\frac{1}{5} \times \frac{1}{4} =$
- Calculate: $p^2 + q^2 + 2pq$ for $p = \frac{1}{3}$ and $q = \frac{2}{3}$
 Also for: $p = \frac{2}{5}$ en $q = \frac{3}{5}$
- Calculate: $p^2 - q^2$ for $p = \frac{3}{4}$ and $q = \frac{1}{4}$
 Also for: $p = \frac{5}{8}$ and $q = \frac{3}{8}$

Multiplying fractions (II)



Mediants (I)

You have seen that:

$$\frac{m}{n} \times \frac{p}{q} = \frac{m \times p}{n \times q} \dots\dots\dots (A)$$

Maybe you should think that:

$$\frac{m}{n} + \frac{p}{q} \stackrel{?}{=} \frac{m+p}{n+q} \dots\dots\dots (B)$$

- Calculate in this way the 'sum' of $\frac{3}{5}$ and $\frac{1}{4}$
Check that this result is lying *between* $\frac{3}{5}$ and $\frac{1}{4}$
so it can not be the real sum of both fractions!
- How can you calculate the right sum of $\frac{3}{5}$ and $\frac{1}{4}$?

Formula (B) is not a good recipe to add fractions,
but it is a recipe to find intermediate fractions, so-called *mediants*.

- Check for some examples, that the values of $\frac{m+p}{n+q}$ always lies between the values of $\frac{m}{n}$ and $\frac{p}{q}$
- Design some examples, in which the value of $\frac{m+p}{n+q}$ lies in the middle between the values of $\frac{m}{n}$ and $\frac{p}{q}$

In a group of 31 students are much more girls than boys
(19 to 12).

In a parallel group (29 students) it is just the reverse (12 to 17)


The two groups are joined during the lessons of P.E.
and then the girls and boys are separated.

- Check that this junction results in a more balanced distribution of numbers of girls and boys.
- What has this example to do with the concept of mediant?

Mediants (II)

With the formula for mediants, can be made step by step new fractions.

- Fill the empty cells in the 'tree' below.



Adding fractions

The correct formula for adding fractions is (unfortunately) much more complicated than the mediant formula. Here it is:

$$\frac{m}{n} + \frac{p}{q} = \frac{m \times q + n \times p}{n \times q}$$

- Check this formula for the fractions $\frac{3}{5}$ and $\frac{1}{4}$
- Design some examples by yourself to check the addition formula.
- Which (simpler) formula do you get if $m = 1$ and $p = 1$?
- Which (insipid) formula do you get if $n = 1$ en $q = 1$?

The addition formula is in many cases more complicated than necessary, but it's always correct.

Here is an example, in which you can add fractions without any formula:

$$\frac{3}{n} + \frac{2}{n} = \frac{5}{n}$$



- If you should use the addition formula (with $m = 3$, $p = 2$ en $q = n$) you should have

$$\frac{3}{n} + \frac{2}{n} = \frac{3 \times n + n \times 2}{n \times n}$$



Show that the result is equal to $\frac{5}{n}$

- Find one fraction equal to: $\frac{m}{3} + \frac{m}{2}$
- Also for: $\frac{m}{3} + \frac{p}{2}$

Trees with fractions (I)



Trees with fractions (II)



More fractions

$$\frac{a}{2} + \frac{a}{3} + \frac{a}{6} = \dots$$

$$\frac{2}{a} + \frac{3}{a} + \frac{6}{a} = \dots$$

$$\frac{b}{2} + \frac{b}{4} + \frac{b}{6} + \frac{b}{12} = \dots$$


$$\frac{2}{b} + \frac{4}{b} + \frac{6}{b} + \frac{12}{b} = \dots$$

$$\frac{c}{2} + \frac{c}{6} + \frac{c}{10} + \frac{c}{12} + \frac{c}{15} + \frac{c}{60} = \dots$$

$$\frac{2}{c} + \frac{6}{c} + \frac{10}{c} + \frac{12}{c} + \frac{15}{c} + \frac{60}{c} = \dots$$

- Design one pair of additions in the same style.

Painting a cube (I)



One cube ($12 \times 12 \times 12$ cm) is glued from little wooden cubes, of 1 cm^3 .

- How many small cubes are used?


The faces of the big cube are painted red.
There are small cubes that get 3 red faces.

- How many?
- How many small cubes get only 2 red faces?
- How many get only 1 red face?
- How many small cubes are not painted at all?

After an idea of Pierre van Hiele.

Painting a cube (II)

A sequence of painted cubes:



The first cube is $3 \times 3 \times 3$ cm, the second one $4 \times 4 \times 4$ cm, etc.

Look at the table below; r represents the length of an edge in cm.

You can read the value of r on the labels in the drawing.



The number of small cubes with 3 painted faces, is called A_3

A_2 , A_1 and A_0 represent the number of small cubes with 2, 1 and 0 painted faces. A_{tot} represents the total number of small cubes.

r	A_3	A_2	A_1	A_0	A_{tot}
3	8	12	6	1	27
4					
5					
6					
7					
8					

- Check the first row in the table and complete the table.
- Which regularities do you notice?
- Explain the formula: $A_2 = 12(r - 2)$
- Try to find expressions in r for A_1 , A_0 and A_{tot}

Triangular numbers and squares





$$\frac{1}{2}n(n-1) + \frac{1}{2}n(n+1) = n^2$$

● Explain :

$$1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9 + 10 + 9 + 8 + 7 + 6 + 5 + 4 + 3 + 2 + 1 = 100$$

Oblong numbers and squares



$$4n(n + 1) + 1 = (2n + 1)^2$$

A remarkable identity (I)

A rectangular field has a length of 31 m and a width of 29 m.

- Give a fast estimation (in m^2) of the area.
- How many m^2 does this estimation deviate from the exact area?

- The same two questions for a field of 41×39 meters

- 51×49
 - estimation: $50 \times 50 = \dots\dots\dots$
 - exact result = $\dots\dots\dots$

deviation = $\dots\dots$


- Make a similar scheme for 61×59 .

- Explain from the picture:

The difference between $n \times n$ and $(n + 1) \times (n - 1)$ is equal to 1

or as a formula:

$$(n + 1) \times (n - 1) = n^2 - 1$$




A remarkable identity (II)

● 32×28 $\left\{ \begin{array}{l} \text{estimation: } 30 \times 30 = \dots\dots\dots \\ \text{exact result} = \dots\dots\dots \end{array} \right. \text{deviation} = \dots\dots$


● Make a similar scheme for 42×38 .

● Also for 52×48 .

● Find a general rule.



A remarkable identity (III)



- Look at the number line and fill:

$$(10 + p) + (10 - p) = \dots\dots\dots$$

$$(10 + p) - (10 - p) = \dots\dots\dots$$

August thinks that ' $10 + p$ times $10 - p$ ' is equal to 100.


His reasoning: **10 times 10 is 100**

$10 - p$ is p less than 10,


but $10 + p$ is p more than 10,

so they compensate each other.

Is August right?



A remarkable identity (IV)



$$a^2 \text{ minus } b^2 = (a + b) \text{ times } (a - b)$$

example:

$$54^2 - 46^2 = (54 + 46) \times (54 - 46) = 100 \times 8 = 800$$

- Calculate in this manner, without calculator:


$$52^2 - 48^2 = \dots = \dots = \dots$$

$$67^2 - 33^2 = \dots = \dots = \dots$$

$$501^2 - 499^2 = \dots = \dots = \dots$$

- Design some exercises in the same style.
- **100** times **100** is more than **100 + p** times **100 - p**
How many more?
- **n²** is more than **n + 10** times **n - 10**
How many more?

Divisible by 6



- Fill in the empty cells.

Look at the strip on the right side. It seems to be that all numbers in that strip are divisible by 6.

- Take some other values for n and investigate if this is also true in those cases.

In the strip with label $n^3 - n$ one can find a nice pattern.



To begin with 6: $6 = 1 \times 2 \times 3$

And then: $24 = 2 \times 3 \times 4$

$60 = 3 \times 4 \times 5$

- Will this be continued? Try some cases.
- Complete (and check!) : $n^3 - n = \dots \times \dots \times \dots$
- Try to explain why $n^3 - n$ is divisible by 6, for each value of n .


Multiplying differences (I)




$$(a - b) \times (c - d) = a \times c - a \times d - b \times c + b \times d$$

- Explain this formula from the pictures
- Is the formula true if $a = b$? Explain your answer.

Multiplying differences (II)



$$(8 - p) \times (5 - q) = \dots 40 \dots$$




$$(x - 3) \times (y - 2) = \dots$$

- Design your own product of differences with corresponding picture.

Multiplying differences (III)

A sheet of paper, size A4, has dimensions 210 and 297 mm.
Rounded to centimeters: 21 and 30 cm.

- Take a sheet A4 and fold it twice: one crease parallel with the shortest edge and one parallel with the other edge.
The distances from the edges are chosen arbitrary.




Call the distance from the 'vertical' crease to the left side L
and the distance from the 'horizontal' crease to the upper side U .

- Unfold the paper; now you see four rectangles.

Write in each of those rectangles an expression in L and U for
the area of that rectangle.

- What will be the result if you add the four expressions?
Check this.

Squares of sums (I)



$$(a + b)^2 = a^2 + b^2 + 2ab$$

- Explain the equivalence of $(a + b)^2$ and $a^2 + b^2 + 2ab$ from the picture.
- Mental arithmetic:
calculate $a^2 + b^2 + 2ab$ in the case that $a = 37$ and $b = 63$
- Suppose $b = a$. The equivalence of both expressions must stay valid..
Verify without using a picture that $(a + a)^2$ and $a^2 + a^2 + 2aa$ are equivalent indeed.
- Suppose $b = 2a$ and check the equivalence.
- Substitute $a = 2p$ and $b = 3p$ in $(a + b)^2$ and $a^2 + b^2 + 2ab$
Verify the equivalence.

Squares of sums (II)

The next poem (translated from Dutch) is from the book "A lark above a pasture" of the author K. Schippers.

The title of this poem is a complicated algebraic formula.

The poem finishes with another formula which looks nicer.

In this way the poet will express the beauty of his lover.

$$a^2 + b^2 + c^2 + 2ab + 2ac + 2bc$$



I say your beauty minus your eyes is a
 the spirit that darts in you is b
 your eyes
 c
 added and at least given a square:
 $(a + b + c)^2$

- The last nice expression is equivalent with the complicated first one. Explain this from the picture.

	a	b	c
a	a^2	ab	ac
b	ab	b^2	bc
c	ac	bc	c^2

- Substitute $c = 0$ in both expressions. Which formula do you get?
- Suppose $a = b = c$. Both expressions can be written using only one letter. Check by calculating that they are equivalent..
- The square of 111 has a funny result: 12321. You can check this by a calculator, but you can also use the two expressions from the poem



Squares of differences (I)



$$(a - b)^2 = a^2 - 2ab + b^2$$

- Explain this formula from the pictures.
- Is the formula correct for $a = b$? Explain your answer.


Squares of differences (II)



$(n - 12)^2 = \dots\dots\dots$

- Design your own square of a difference and draw the corresponding picture.

More about squares



$$(a - b)^2 + (a + b)^2 = 2a^2 + 2b^2$$

- Explain this formula.
- Mental arithmetic: $99^2 + 101^2$
- Also: $49^2 + 50^2 + 51^2$

$$S = (n - 2)^2 + (n - 1)^2 + n^2 + (n + 1)^2 + (n + 2)^2$$


Assertion: S is divisible by 5, for each integer value of n

- Investigate if this assertion is true. If it's true, try to give an explanation.

$$T = (n - 3)^2 + (n - 2)^2 + (n - 1)^2 + n^2 + (n + 1)^2 + (n + 2)^2 + (n + 3)^2$$

- Which number will divide T , regardless the value of n ?
Give an explanation.

Sum, difference, product



$$(a + b)^2 - (a - b)^2 = 4ab$$

- Explain this formula from the picture.
- Check the formula in the case: $a = b$.

$$S^2 - D^2 = (S + D)(S - D)$$


- Suppose $S = a + b$ and $D = a - b$
Substitute this in the preceding formula.
This leads to the formula in the black rectangle.

S represents the sum, **D** the difference and **P** the product of two numbers.

- Complete this table:


S	D	P
13	3	
22		96
	19	120
	0	576

Equations with squares (I)




area = 441
or:
 $(x + 6)^2 = 441$

- Calculate the value of x.



area = 400
or:
 $(\dots\dots\dots)^2 = 400$


- Calculate the value of y.



area = 364
or:
 $z^2 + 4 \times 3z = 364$

- Calculate the value of z.

Equations with squares (II)




area = 396

or:

..... = 396

- Calculate the value of a .





area = 384

or:



..... = 384

- Calculate the value of b .

Equations with squares (III)



- Calculate the value of p .



- Calculate the value of x .

Antique equation (I)

The square of the with 3 reduced fifth part of a group of monkeys was hidden in a cave.

Only one monkey, who climbed in a tree, was visible.

How many monkeys were there totally?

Bhaskara, India (1114 - 1185)

Antique equation (II)


*The son of Pritha, rancorous by the fight,
used a quiver full of arrows
to kill Karna:*

- with half of his arrows he repelled
the arrows of his enemy;*
- with 4 times the square root of the total
number of arrows he killed his horse;*
- with 6 arrows he brought down Salya;*
- with 3 he destroyed his sunshade and bow*
- with 1 arrow he hitted the head of the fool.*

*How many arrows were shoot by Arjuna,
the son of Pritha?*

Bhaskara, from Vya Ganita (= 'calculus of square roots')

Factorizing (I)



- How can you make one rectangle with the four pieces?
- Explain: $xy + x + y + 1 = (x + 1)(y + 1)$
- Complete: $xy + 2x + 2y + 4 = (x + \dots)(y + \dots)$
How can you explain the equivalence by a picture?
- Fill in the missing numbers:

$$ab + 2a + 3b + 6 = (a + \dots)(b + \dots)$$

$$pq + 5p + 6q + 30 = (p + \dots)(q + \dots)$$

$$xy + 11x + 9y + \dots = (x + 9)(y + \dots)$$

$$mn + \dots m + \dots n + 200 = (m + 20)(n + \dots)$$

- Suppose $b = a$, $q = p$, $y = x$ and $n = m$, then you get (fill in):


$$a^2 + 5a + 6 = (a + \dots)(a + \dots)$$

$$p^2 + 11p + 30 = (p + \dots)(p + \dots)$$

$$x^2 + 20x + \dots = (x + 9)(x + \dots)$$

$$m^2 + \dots m + 200 = (m + 20)(m + \dots)$$

Factorizing (II)



- Check the calculation above.
- Make a similar calculation, starting with respectively:
 $x + 4$, $y + 10$, $z + 11$, $p + 1$
- Create some other examples yourself.
- Try to give a general rule?


Formula with cubes(I)

In an old algebra book one can find this formula:

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$

Without a real explanation, you can carry out some tests to investigate if the formula could be correct.

- Exchange a and b .



- Suppose $b = 0$

Then: $(a + b)^3 = \dots\dots\dots$ and $a^3 + 3a^2b + 3ab^2 + b^3 = \dots\dots\dots$

- Suppose $b = a$

Then: $(a + b)^3 = \dots\dots\dots$ and $a^3 + 3a^2b + 3ab^2 + b^3 = \dots\dots\dots$


- Suppose $b = 10a$

Then: $(a + b)^3 = \dots\dots\dots$ and $a^3 + 3a^2b + 3ab^2 + b^3 = \dots\dots\dots$

All these tests don't give any guarantee that the formula is correct, they only give some trust.

Look at the eight blocks.

- How can they be used to explain the formula at the top of this page?




Formula with cubes (II)


$$(a + b + c)^3 = a^3 + 3a^2b + \dots + c^3$$

- Complete this formula.

Hint: look at the picture and imagine that it will be extended to a cube building with three floors, respectively with height **a**, **b** and **c**. On each floor are nine 'rooms'. You can find an expression for the volume of each room.....



A remarkable pattern



- Check that the the four results are squares.
- How should you continue the list? Give two lines more.
Check that the results are squares again.

The product of any four consecutive whole numbers added to 1 is a square!

- Try to prove this rule by using algebra.
Here are two hints:
 - * compare the product of the two inner factors with the product of the two outer ones;
 - * look for a relationship between these products and the base of the square.